

Cosmology in massive gravity

Yungui Gong^{1,2}

¹*School of Physics, Huazhong University of Science and Technology, Wuhan 430074, China*

²*Institute of Theoretical Physics, Chinese Academy of Sciences, Beijing 100190, China**

We argue that more cosmological solutions in massive gravity can be obtained if the metric tensor and the tensor $\Sigma_{\mu\nu}$ defined by Stückelberg fields take the homogeneous and isotropic form. The standard cosmology with matter and radiation dominations in the past can be recovered and Λ CDM model is easily obtained. The dynamical evolution of the universe is modified at very early times.

PACS numbers: 04.50.Kd, 98.80.-k

The discovery of accelerating expansion of the universe by the observations of Type Ia supernovae in 1998 [1, 2] motivated the search for dark energy and modified gravity. Dvali, Gabadadze and Porrati proposed that general relativity is modified at the cosmological scale [3]. It is also possible that graviton has a small mass which is comparable to the Hubble scale. The first attempt of a theory of gravity with massive graviton was made by Fierz and Pauli [4]. However, the linear theory with the Fierz-Pauli mass is in contradiction with solar system tests [5, 6]. Recently, de Rham, Gabadadze and Tolley introduced a nonlinear theory of massive gravity [7] that is free from Boulware-Deser ghost [8, 9]. Cosmological solutions for massive gravity were sought in [10–16]. The first homogenous and isotropic solution was found for spatially open universe in [11] and the massive graviton term is equivalent to a cosmological constant. The same solutions were then found for spatially open and closed universe in [12, 13]. Langlois and Naruko then found more general cosmological solutions [16]. We follow the approach in [16] and proposed a new approach to get more general cosmological solutions.

The theory of massive gravity is base on the following action [7]

$$S = \frac{M_{pl}^2}{2} \int d^4x \sqrt{-g} (R + m_g^2 \mathcal{U}) + S_m, \quad (1)$$

where m_g is the mass of the graviton, the mass term

$$\mathcal{U} = \mathcal{U}_2 + \alpha_3 \mathcal{U}_3 + \alpha_4 \mathcal{U}_4, \quad (2)$$

$$\mathcal{U}_2 = [\mathcal{K}]^2 - [\mathcal{K}^2], \quad (3)$$

$$\mathcal{U}_3 = [\mathcal{K}]^3 - 3[\mathcal{K}][\mathcal{K}^2] + 2[\mathcal{K}^3], \quad (4)$$

$$\mathcal{U}_4 = [\mathcal{K}]^4 - 6[\mathcal{K}]^2[\mathcal{K}^2] + 8[\mathcal{K}^3][\mathcal{K}] - 6[\mathcal{K}^4], \quad (5)$$

and

$$\mathcal{K}_\nu^\mu = \delta_\nu^\mu - (\sqrt{\Sigma})_\nu^\mu, \quad (6)$$

The tensor $\Sigma_{\mu\nu}$ is defined by four Stückelberg fields ϕ^a as

$$\Sigma_{\mu\nu} = \partial_\mu \phi^a \partial_\nu \phi^b \eta_{ab}. \quad (7)$$

The reference metric η_{ab} is usually taken as the Minkowski one. The cosmological solution was first found in [11] for open universe and the mass term behaves like an effective cosmological constant with

$$\Lambda_{eff} = -m_g^2 \left(1 + 3\alpha_3 \pm \sqrt{1 + 3\alpha_3 + 9\alpha_3^2 - 12\alpha_4} \right) \times \frac{(1 + 9\alpha_3^2 - 24\alpha_4 \pm (1 + 3\alpha_3)\sqrt{1 + 3\alpha_3 + 9\alpha_3^2 - 12\alpha_4})}{9(\alpha_3 + 4\alpha_4)^2}. \quad (8)$$

The same solution was then found in [12] for flat universe. For a general case with positive, negative and zero curvature, Kobayashi *et al.* found the same solution with $\Lambda_{eff} = m_g^2/\alpha$ for the particular choices of parameters α_3 and α_4 [13],

$$\alpha_3 = \frac{1}{3}(\alpha - 1), \quad \alpha_4 = \frac{1}{12}(\alpha^2 - \alpha + 1). \quad (9)$$

In [16], the authors assumed that the spacetime metric takes the form

$$ds^2 = g_{\mu\nu} dx^\mu dx^\nu = -N^2(t) dt^2 + a^2(t) \gamma_{ij}(x) dx^i dx^j, \quad (10)$$

with the spatial metric

$$\gamma_{ij}(x) dx^i dx^j = \frac{dr^2}{1 - kr^2} + r^2(d\theta^2 + \sin^2 \theta d\phi^2),$$

and generalized the reference metric from Minkowski metric to de Sitter metric,

$$\eta_{ab} d\phi^a d\phi^b = -dT^2 + b_k^2(T) \gamma_{ij} dX^i dX^j, \quad (11)$$

where the Stückelberg fields are assumed to be $\phi^0 = T = f(t)$, $\phi^i = X^i = x^i$, so that the tensor $\Sigma_{\mu\nu}$ takes the homogeneous and isotropic form,

$$\Sigma_{\mu\nu} = \text{Diag}\{-\dot{f}^2, b_k^2[f(t)]\gamma_{ij}\}, \quad (12)$$

and the functions $b_k(T)$ are

$$b_0(T) = e^{H_c T}, \quad b_{-1}(T) = H_c^{-1} \sinh(H_c T),$$

$$b_1(T) = H_c^{-1} \cosh(H_c T),$$

they then found three branches of cosmological solutions, two of them correspond to the effective cosmological constant (8) and exist for spatially flat, open and closed cases. They also found a new solution

$$\frac{db_k[f]}{df} = \frac{\dot{a}}{N}. \quad (13)$$

For the flat case, $k = 0$, substituting the de Sitter function $b_0[f(t)] = e^{H_c f(t)}$ into equation (13), we obtain the effective energy density and pressure for the massive graviton,

$$\rho_g = -m_g^2 M_{pl}^2 \left(1 - \frac{H}{H_c}\right) \left[3(\alpha_3 + 4\alpha_4) \frac{H^2}{H_c^2} - 3(1 + 5\alpha_3 + 8\alpha_4) \frac{H}{H_c} + 6 + 12\alpha_3 + 12\alpha_4\right], \quad (14)$$

$$p_g = m_g^2 M_{pl}^2 \left[-3(\alpha_3 + 4\alpha_4) \frac{H^3}{H_c^3} \left(1 + \frac{\dot{H}}{H^2}\right) + 6 + 12\alpha_3 + 12\alpha_4 - (3 + 9\alpha_3 + 12\alpha_4) \frac{H}{H_c} \left(3 + \frac{\dot{H}}{H^2}\right) + (1 + 6\alpha_3 + 12\alpha_4) \frac{H^2}{H_c^2} \left(3 + 2\frac{\dot{H}}{H^2}\right) \right]. \quad (15)$$

So when $H = H_c$, $\rho_g = 0$. The Friedmann equations are

$$H^2 + \frac{k}{a^2} = \frac{1}{3M_{pl}^2}(\rho_m + \rho_g), \quad (16)$$

$$2\dot{H} + 3H^2 + \frac{k}{a^2} = -\frac{1}{M_{pl}^2}(p_m + p_g). \quad (17)$$

For the flat case, substituting equations (14) and (15) into Friedmann equations (16) and (17), we get

$$E^2(z) + \frac{m_g^2}{H_0^2} \frac{H(z)}{H_c} \left[-(\alpha_3 + 4\alpha_4) \frac{H^2(z)}{H_c^2} + (1 + 6\alpha_3 + 12\alpha_4) \frac{H(z)}{H_c} - 3(1 + 3\alpha_3 + 4\alpha_4) \right] = \Omega_m(1+z)^{3(1+w_m)} - 2\frac{m_g^2}{H_0^2}(1 + 2\alpha_3 + 2\alpha_4). \quad (18)$$

$$\frac{\dot{H}}{H^2} \left\{ \left[3(1 + 3\alpha_3 + 4\alpha_4) \frac{H_c}{H_0} - 2(1 + 6\alpha_3 + 12\alpha_4) E(z) + 3(\alpha_3 + 4\alpha_4) \frac{H_0}{H_c} E^2(z) \right] \frac{m_g^2}{H_c^2} E(z) - 2E^2(z) \right\} = 3\Omega_m(1+w_m)(1+z)^{3(1+w_m)}, \quad (19)$$

where $E(z) = H(z)/H_0$. The effective equation of state $w_g = p_g/\rho_g$ for the massive graviton is

$$w_g = -1 - \frac{2E^2(z)\dot{H}/H^2 + 3\Omega_m(1+w)(1+z)^{3(1+w_m)}}{3[E^2(z) - \Omega_m(1+z)^{3(1+w_m)}]}. \quad (20)$$

Since $\rho_g = 0$ when $H(z) = H_c$, so if $H_c = H_0$, we find that $\Omega_m = 1$ which is inconsistent with current observations, therefore $H_c \neq H_0$. If $H_c < H_0$, then we cannot recover the standard cosmology $H^2 \sim \rho$ in the past unless we fine tune the value of m_g^2/H_0^2 to be very small. From equation (18), we see that the standard cosmology is recovered when $H(z) \ll H_c$ and $m_g \ll H_0 < H_c$. At very early times, $H(z) > H_c$, the universe evolves according to $H^3 \sim \rho$. If it was radiation dominated in the very early times, then the universe evolves faster as $a(t) \sim t^{3/4}$ instead of $t^{1/2}$. For the special case $\alpha_3 = \alpha_4 = 0$, Friedmann equation is simplified to

$$\left(1 + \frac{m_g^2}{H_c^2}\right) E^2(z) - 3\frac{m_g^2}{H_c H_0} E(z) + 2\frac{m_g^2}{H_0^2} = \Omega_m(1+z)^{3(1+w_m)}. \quad (21)$$

At $z = 0$, $E(z) = 1$, we get

$$\frac{m_g^2}{H_0^2} = -\frac{1 - \Omega_m}{(H_0/H_c - 2)(H_0/H_c - 1)}. \quad (22)$$

As discussed above, $H_c/H_0 > 1$, so m_g^2 must be negative for this special case. Without loss of generality, we assume that $m_g^2 = -\beta_1 H_0^2$, and $H_c = \beta_2 H_0$ with $\beta_2 > 1$. For the special case $\alpha_3 = \alpha_4 = 0$, equation (22) gives

$$\beta_1 = \frac{(1 - \Omega_m)\beta_2^2}{(2\beta_2 - 1)(\beta_2 - 1)}. \quad (23)$$

In this case, we have only two free parameters Ω_m and β_2 , and equation (19) gives

$$\frac{\dot{H}}{H^2} = -\frac{3\Omega_m(1+w_m)(1+z)^{3(1+w_m)}}{2\left(1 - \frac{\beta_1}{\beta_2^2}\right) E^2(z) + 3\frac{\beta_1}{\beta_2} E(z)}. \quad (24)$$

If $\beta_2 \gg 1$, then $\beta_1 = (1 - \Omega_m)/2$, and the model becomes the Λ CDM model. This is shown in Fig. 1 for $\beta_2 = 20.1$ and $\Omega = 0.3$.

In summary, starting with the homogeneous and isotropic metric (10) and tensor (12), we obtain the Friedmann equations (16) and (17) with the effective energy density and pressure for the massive graviton,

$$\rho_g = \frac{m_g^2 M_{pl}^2}{a^3} (b_k[f] - a) \{6(1 + 2\alpha_3 + 2\alpha_4) a^2 - (3 + 15\alpha_3 + 24\alpha_4) a b_k[f] + 3(\alpha_3 + 4\alpha_4) b_k[f]^2\}, \quad (25)$$

$$p_g = \frac{m_g^2 M_{pl}^2}{a^2} \{ [6 + 12\alpha_3 + 12\alpha_4 - (3 + 9\alpha_3 + 12\alpha_4) \dot{f}] a^2 - 2[3 + 9\alpha_3 + 12\alpha_4 - (1 + 6\alpha_3 + 12\alpha_4) \dot{f}] a b_k[f] + [1 + 6\alpha_3 + 12\alpha_4 - 3(\alpha_3 + 4\alpha_4) \dot{f}] b_k[f]^2 \}, \quad (26)$$

and the equation of motion for the function $f(t)$ which




FIG. 1: The evolution of the deceleration parameter $q(z)$ and the effective equation of state w_g of massive graviton. The blue lines are for the model with de Sitter metric as the reference metric and the black lines are for the model taking $b_k(f)$ as power law form.

leads to the three branches of solutions

$$b_k[f(t)] = \frac{(1 + 6\alpha_3 + 12\alpha_4 \pm \sqrt{1 + 3\alpha_3 + 9\alpha_3^2 - 12\alpha_4})}{3(\alpha_3 + 4\alpha_4)} a(t), \quad (27)$$

$$\frac{db_k[f]}{df} = \frac{\dot{a}}{N}. \quad (28)$$

When we take the solution (27), we get the effective cosmological constant solution (8) independent of the choice of spatial curvature k . For $k = 1$, the solution was obtained in [11] by taking the reference metric η_{ab} as Minkowski and the same homogeneous and isotropic tensor $\Sigma_{\mu\nu}$ (12) with $b_k[f(t)] = f(t)$. The same solution (8)

was obtained in [13] for all values of k for the particular parameters (9), but the tensor $\Sigma_{\mu\nu}$ is not homogeneous and isotropic. For the flat case $k = 0$, Gratia, Hu and Wyman obtained the same cosmological constant solution (8) by using another inhomogeneous and anisotropic tensor $\Sigma_{\mu\nu}$ [12]. However, the cosmological constant solution (8) is just the consequence of the equation of motion of the function $f(t)$ once we assumed the homogeneous and isotropic form for the metric (10) and the tensor $\Sigma_{\mu\nu}$ (12). This suggests that the solution (28) is quite general even without the assumption of η_{ab} as de Sitter. Therefore, we propose that more solutions can be found from this approach by assuming more general form of $b_k[f(t)]$.

Follow the above argument, we assume a power law form $b_k[f(t)] = (H_c f(t))^{\gamma/(\gamma-1)}$ with $\gamma > 1$, then the solution to equation (28) is

$$b_k[f(t)] = \left(\frac{a(\gamma-1)H}{\gamma H_c} \right)^\gamma, \quad (29)$$

and the effective energy density becomes

$$\begin{aligned} \rho_g = 3m_g^2 M_{pl}^2 & \left[3(1 + 3\alpha_3 + 4\alpha_4) \left(\frac{\gamma-1}{\gamma} \right)^\gamma a^{\gamma-1} \left(\frac{H}{H_c} \right)^\gamma \right. \\ & - (1 + 6\alpha_3 + 12\alpha_4) \left(\frac{\gamma-1}{\gamma} \right)^{2\gamma} a^{2\gamma-2} \left(\frac{H}{H_c} \right)^{2\gamma} \\ & - 2(1 + 2\alpha_3 + 2\alpha_4) \\ & \left. + (\alpha_3 + 4\alpha_4) \left(\frac{\gamma-1}{\gamma} \right)^{3\gamma} a^{3\gamma-3} \left(\frac{H}{H_c} \right)^{3\gamma} \right]. \end{aligned} \quad (30)$$

The effective pressure of massive graviton is

$$\begin{aligned} p_g = m_g^2 M_{pl}^2 & \left\{ 6(1 + 2\alpha_3 + 2\alpha_4) - 3(1 + 3\alpha_3 + 4\alpha_4) \left(\frac{\gamma-1}{\gamma} \right)^\gamma a^{\gamma-1} \left(\frac{H}{H_c} \right)^\gamma \left[2 + \gamma \left(1 + \frac{\dot{H}}{H^2} \right) \right] \right. \\ & + (1 + 6\alpha_3 + 12\alpha_4) \left(\frac{\gamma-1}{\gamma} \right)^{2\gamma} a^{2\gamma-2} \left(\frac{H}{H_c} \right)^{2\gamma} \left[1 + 2\gamma \left(1 + \frac{\dot{H}}{H^2} \right) \right] \\ & \left. - 3\gamma(\alpha_3 + 4\alpha_4) \left(\frac{\gamma-1}{\gamma} \right)^{3\gamma} a^{3\gamma-3} \left(\frac{H}{H_c} \right)^{3\gamma} \left(1 + \frac{\dot{H}}{H^2} \right) \right\}. \end{aligned} \quad (31)$$

Again when the hubble parameter is in the range $H_0 < H(z) < H_c$, the standard cosmology is recovered. To have a long history of matter and radiation domination,

we require $H_0 \ll H_c$. For the special case $\alpha_3 = \alpha_4 = 0$, Friedmann equations are

$$E^2(z) - \frac{\beta_1}{\beta_2^{2\gamma}} \left(\frac{\gamma-1}{\gamma} \right)^{2\gamma} a^{2\gamma-2} E^{2\gamma}(z) - 2\beta_1 + 3\frac{\beta_1}{\beta_2^\gamma} \left(\frac{\gamma-1}{\gamma} \right)^\gamma a^{\gamma-1} E^\gamma(z) = \Omega_m(1+z)^{3(1+w_m)}, \quad (32)$$

$$\frac{\dot{H}}{H} = -1 + \frac{\frac{\beta_1}{\beta_2^{2\gamma}} \left(\frac{\gamma-1}{\gamma}\right)^{2\gamma} a^{2\gamma-2} E^{2\gamma} - 6 \frac{\beta_1}{\beta_2^2} \left(\frac{\gamma-1}{\gamma}\right)^\gamma a^{\gamma-1} E^\gamma - E^2 + 6\beta_1}{-2\gamma \frac{\beta_1}{\beta_2^{2\gamma}} \left(\frac{\gamma-1}{\gamma}\right)^{2\gamma} a^{2\gamma-2} E^{2\gamma} + 3\gamma \frac{\beta_1}{\beta_2^2} \left(\frac{\gamma-1}{\gamma}\right)^\gamma a^{\gamma-1} E^\gamma + 2E^2}, \quad (33)$$

with

$$\beta_1 = \frac{(1 - \Omega_m) \beta_2^{2\gamma}}{\left[\left(\frac{\gamma-1}{\gamma}\right)^\gamma - \beta_2^\gamma\right] \left[\left(\frac{\gamma-1}{\gamma}\right)^\gamma - 2\beta_2^\gamma\right]}. \quad (34)$$

In this case, we have three free parameters Ω_m , γ and β_2 . Again if $\beta_2 \gg 1$, $\beta_1 = (1 - \Omega_m)/2$ and the model is equivalent to the Λ CDM model. This is shown in Fig. 1 for $\beta_2 = 20.1$ and $\Omega = 0.3$.

In conclusion, more general cosmological solutions which are consistent with observational data can be found by taking homogeneous and isotropic form for both the metric $g_{\mu\nu}$ and the tensor $\Sigma_{\mu\nu}$. The mass of graviton is in the order of $((1 - \Omega_m)/2)^{1/2} H_0$.

This work was partially supported by the National Basic Science Program (Project 973) of China under grant Nos. 2007CB815401 and 2010CB833004, the NNSF of China under grant Nos. 10935013 and 11175270, CQ CSTC under grant No. 2009BA4050 and CQ CMEC under grant No. KJTD201016.

-
- [1] A.G. Riess *et al.*, *Astron. J.* **116**, 1009 (1998).
 - [2] S. Perlmutter *et al.*, *Astrophys. J.* **517**, 565 (1999).
 - [3] G. Dvali, G. Gabadadze and M. Porrati, *Phys. Lett. B* **485**, 208 (2000).
 - [4] M. Fierz and W. Pauli, *Proc. Roy. Soc. Lond. A* **173**, 211 (1939).
 - [5] H. van Dam and M. Veltman, *Nucl. Phys. B* **22**, 397 (1970).
 - [6] V. Zakharov, *JETP Lett.* **12**, 312 (1970).
 - [7] C. de Rham, G. Gabadadze and A.J. Tolley, *Phys. Rev. Lett.* **106**, 231101 (2011).
 - [8] D. Boulware and S. Deser, *Phys. Lett. B* **40**, 227 (1972).
 - [9] S. Hassan and R.A. Rosen, *Phys. Rev. Lett.* **108**, 041101 (2012).
 - [10] G. D'Amico *et al.*, *Phys. Rev. D* **84**, 124046 (2011).
 - [11] A.E. Gumrukcuoglu, C. Lin and S. Mukohyama, *J. Cosm. Astropart. Phys.*, JCAP11(2011)030.
 - [12] T. Gratia, W. Hu and M. Wyman, arXiv: 1205.4241.
 - [13] T. Kobayashi, M. Siino, M. Yamaguchi, and D. Yoshida, arXiv: 1205.4838.
 - [14] M.S. Volkov, arXiv: 1205.5713.
 - [15] A.E. Gümrükçüoğlu, C. Lin and S. Mukohyama, arXiv: 1206.2723.
 - [16] D. Langlois and A. Naruko, arXiv: 1206.6810.

* Electronic address: ygong@mail.hust.edu.cn